

Original article

- 1 Terashima M, Ikeda K, Maesawa C, Kawamura H, Niitsu Y, Satoh M, Saito K. Establishment of an alpha-fetoprotein-producing gastric cancer cell line in serum-free media. *Jpn J Cancer Res* 1991;82:883-885.
- 2 Tamura G, Maesawa C, Mikawa S, Satodate R. Significance of immunohistochemically detected c-erbB-2 protein expression in stage III breast cancer, with reference to nuclear deformity, DNA content and prognosis. *Jpn J Clin Oncol* 1991;21:264-267.
3. Terashima M, Ikeda K, Kawamura S, Maesawa C, Ishida K, Sato M, Saito K. The usefulness of cyclophosphamide pretreatment for subrenal capsule assay against human esophageal cancer. *Jpn J Surg* 1991;21:184-192.
- 4 Maesawa C, Tamura G, Suzuki Y, Ishida K, Saito K, Satodate R. Sensitive detection of p53 gene mutations in esophageal endoscopic biopsy specimens by cell sorting combined with polymerase chain reaction single-strand conformation polymorphism analysis. *Jpn J Cancer Res* 1992;83:1253-1256.
- 5 Maesawa C, Masuda T, Tamura G, Satodate R, Ishida K, Saito K. Prognostic assessment of superficial squamous cell carcinoma of the esophagus using karyometric analysis and nucleolar organizer regions. *J Surg Oncol* 1992;51:164-168.
- 6 Tamura G, Maesawa C, Suzuki Y, Satodate R, Ishida K, Saito K. p53 gene mutations in esophageal cancer detected by polymerase chain reaction single-strand conformation polymorphism analysis. *Jpn J Cancer Res* 1992;83:559-562.
- 7 ©Maesawa C, Sakuma T. Microscopic features of the regeneration of white pulp in autotransplanted spleens in rats. *Virchows Arch A Pathol Anat Histopathol* 1992;421:41-45.
- 8 Terashima M, Ikeda K, Maesawa C, Kawamura H, Ishida K, Sato M, Saito K. Drug-sensitivity testing in patients with human oesophageal squamous cell carcinoma. *Eur J Cancer* 1992;28A:1347-1350.
- 9 Tamura G, Maesawa C, Suzuki Y, Ogasawara S, Terashima M, Saito K, Satodate R. Primary gastric carcinoma cells frequently lose heterozygosity at the APC and MCC genetic loci. *Jpn J Cancer Res* 1993;84:1015-1018.
- 10 ○Maesawa C, Tamura G, Suzuki Y, Ogasawara S, Ishida K, Saito K, Satodate R. Aberrations of tumor-suppressor genes (p53, apc, mcc and Rb) in esophageal squamous-cell carcinoma. *Int J Cancer* 1994;57:21-25.
- 11 Tamura G, Maesawa C, Suzuki Y, Tamada H, Satoh M, Ogasawara S, Kashiwaba M, Satodate R. Mutations of the APC gene occur during early stages of gastric adenoma development. *Cancer Res* 1994;54:1149-1151.
- 12 Ogasawara S, Maesawa C, Tamura G, Satodate R. Lack of mutations of the adenomatous polyposis coli gene in oesophageal and gastric carcinomas. *Virchows Arch*

- 1994;424:607-611.
- 13 Suzuki Y, Tamura G, Maesawa C, Fujioka T, Kubo T, Satodate R. Analysis of genetic alterations in renal cell carcinoma using the polymerase chain reaction. *Virchows Arch* 1994;424:453-457.
 - 14 Tamura G, Maesawa C, Suzuki Y, Kashiwaba M, Ishida M, Saito K, Satodate R. Improved detection of loss of heterozygosity at retinoblastoma gene locus in human breast carcinoma. *Pathol Int* 1994;44:34-38.
 - 15 Suzuki Y, Tamura G, Maesawa C, Fujioka T, Kubo T, Satodate R. Analysis of genetic alterations in renal cell carcinoma using the polymerase chain reaction. *Virchows Arch* 1994;425:335.
 - 16 Kashiwaba M, Tamura G, Suzuki Y, Maesawa C, Ogasawara S, Sakata K, Satodate R. Epithelial-cadherin gene is not mutated in ductal carcinomas of the breast. *Jpn J Cancer Res* 1995;86:1054-1059.
 - 17 Maesawa C, Sakuma T, Sato T, Masuda T, Muro-oka G, Satodate R. Structural characteristic of splenic sinuses in idiopathic portal hypertension. *Pathol Int* 1995;45:642-648.
 - 18 ◦Maesawa C, Tamura G, Suzuki Y, Ogasawara S, Sakata K, Kashiwaba M, Satodate R. The sequential accumulation of genetic alterations characteristic of the colorectal adenoma-carcinoma sequence does not occur between gastric adenoma and adenocarcinoma. *J Pathol* 1995;176:249-258.
 - 19 Tamura G, Sakata K, Maesawa C, Suzuki Y, Terashima M, Satoh K, Sekiyama S, Suzuki A, Eda Y, Satodate R. Microsatellite alterations in adenoma and differentiated adenocarcinoma of the stomach. *Cancer Res* 1995;55:1933-1936.
 - 20 Sakata K, Tamura G, Maesawa C, Suzuki Y, Terashima M, Satoh K, Eda Y, Suzuki A, Sekiyama S, Satodate R. Loss of heterozygosity on the short arm of chromosome 9 without p16 gene mutation in gastric carcinomas. *Jpn J Cancer Res* 1995;86:333-335.
 - 21 ◦Ogasawara S, Maesawa C, Tamura G, Satodate R. Frequent microsatellite alterations on chromosome 3p in esophageal squamous cell carcinoma. *Cancer Res* 1995;55:891-894.
 - 22 Tamura G, Sakata K, Nishizuka S, Maesawa C, Suzuki Y, Terashima M, Eda Y, Satodate R. Allelotype of adenoma and differentiated adenocarcinoma of the stomach. *J Pathol* 1996;180:371-377.
 - 23 Tamura G, Sakata K, Nishizuka S, Maesawa C, Suzuki Y, Iwaya T, Terashima M, Saito K, Satodate R. Inactivation of the E-cadherin gene in primary gastric carcinomas and gastric carcinoma cell lines. *Jpn J Cancer Res* 1996;87:1153-1159.
 - 24 ◦Maesawa C, Tamura G, Nishizuka S, Ogasawara S, Ishida K, Terashima M, Sakata K, Sato N, Saito K, Satodate R. Inactivation of the CDKN2 gene by homozygous deletion and de novo methylation is associated with advanced stage esophageal squamous cell carcinoma.

- Cancer Res 1996;56:3875-3878.
- 25 Maesawa C, Tamura G, Ogasawara S, Suzuki Y, Sakata K, Sugimura J, Nishizuka S, Sato N, Ishida K, Saito K, Satodate R. Loss of heterozygosity at the DCC gene locus is not crucial for the acquisition of metastatic potential in oesophageal squamous cell carcinoma. *Eur J Cancer* 1996;32A:896-898.
- 26 Tamura G, Ogasawara S, Nishizuka S, Sakata K, Maesawa C, Suzuki Y, Terashima M, Saito K, Satodate R. Two distinct regions of deletion on the long arm of chromosome 5 in differentiated adenocarcinomas of the stomach. *Cancer Res* 1996;56:612-615.
- 27 Ogasawara S, Tamura G, Maesawa C, Suzuki Y, Ishida K, Satoh N, Uesugi N, Saito K, Satodate R. Common deleted region on the long arm of chromosome 5 in esophageal carcinoma. *Gastroenterology* 1996;110:52-57.
- 28 Tamura G, Sakata K, Nishizuka S, Maesawa C, Suzuki Y, Iwaya T, Terashima M, Saito K, Satodate R. Analysis of the fragile histidine triad gene in primary gastric carcinomas and gastric carcinoma cell lines. *Genes Chromosomes Cancer* 1997;20:98-102.
- 29 Maesawa C, Tamura G, Nishizuka S, Iwaya T, Ogasawara S, Ishida K, Sakata K, Sato N, Ikeda K, Kimura Y, Saito K, Satodate R. MAD-related genes on 18q21.1, Smad2 and Smad4, are altered infrequently in esophageal squamous cell carcinoma. *Jpn J Cancer Res* 1997;88:340-343.
- 30 Nishizuka S, Tamura G, Maesawa C, Sakata K, Suzuki Y, Iwaya T, Terashima M, Saito K, Satodate R. Analysis of the DPC4 gene in gastric carcinoma. *Jpn J Cancer Res* 1997;88:335-339.
- 31 Sakata K, Tamura G, Nishizuka S, Maesawa C, Suzuki Y, Iwaya T, Terashima M, Saito K, Satodate R. Commonly deleted regions on the long arm of chromosome 21 in differentiated adenocarcinoma of the stomach. *Genes Chromosomes Cancer* 1997;18:318-321.
- 32 Iwaya T, Maesawa C, Nishizuka S, Suzuki Y, Sakata K, Sato N, Ikeda K, Koeda K, Ogasawara S, Otsuka K, Kimura Y, Aoki K, Ishida K, Saito K, Tamura G. Infrequent frameshift mutations of polynucleotide repeats in multiple primary cancers affecting the esophagus and other organs. *Genes Chromosomes Cancer* 1998;23:317-322.
- 33 Nozawa H, Oda E, Ueda S, Tamura G, Maesawa C, Muto T, Taniguchi T, Tanaka N. Functionally inactivating point mutation in the tumor-suppressor IRF-1 gene identified in human gastric cancer. *Int J Cancer* 1998;77:522-527.
- 34 Ikeda K, Terashima M, Kawamura H, Takiyama I, Koeda K, Takagane A, Sato N, Ishida K, Iwaya T, Maesawa C, Yoshinari H, Saito K. Pharmacokinetics of cisplatin in combined cisplatin and 5-fluorouracil therapy: a comparative study of three different schedules of cisplatin administration. *Jpn J Clin Oncol* 1998;28:168-175.
- 35 Iwaya T, Maesawa C, Ogasawara S, Tamura G. Tylosis esophageal cancer locus on

- chromosome 17q25.1 is commonly deleted in sporadic human esophageal cancer. *Gastroenterology* 1998;114:1206-1210.
- 36 Maesawa C, Tamura G, Iwaya T, Ogasawara S, Ishida K, Sato N, Nishizuka S, Suzuki Y, Ikeda K, Aoki K, Saito K, Satodate R. Mutations in the human homologue of the *Drosophila* patched gene in esophageal squamous cell carcinoma. *Genes Chromosomes Cancer* 1998;21:276-279.
- 37 Saito A, Kanai Y, Maesawa C, Ochiai A, Torii A, Hirohashi S. Disruption of E-cadherin-mediated cell adhesion systems in gastric cancers in young patients. *Jpn J Cancer Res* 1999;90:993-999.
- 38 Tamura G, Nishizuka S, Maesawa C, Suzuki Y, Iwaya T, Sakata K, Endoh Y, Motoyama T. Mutations in mitochondrial control region DNA in gastric tumours of Japanese patients. *Eur J Cancer* 1999;35:316-319.
- 39 Satoh M, Nakamura M, Saitoh H, Satoh H, Maesawa C, Segawa I, Tashiro A, Hiramori K. Tumor necrosis factor-alpha-converting enzyme and tumor necrosis factor-alpha in human dilated cardiomyopathy. *Circulation* 1999;99:3260-3265.
- 40 Yamada T, Takaoka AS, Naishiro Y, Hayashi R, Maruyama K, Maesawa C, Ochiai A, Hirohashi S. Transactivation of the multidrug resistance 1 gene by T-cell factor 4/beta-catenin complex in early colorectal carcinogenesis. *Cancer Res* 2000;60:4761-4766.
- 41 Ikeda K, Ishida K, Sato N, Koeda K, Aoki K, Kimura Y, Iwaya T, Ogasawara S, Iijima S, Nakamura R, Uesugi N, Maesawa C, Saito K. Chemoradiotherapy followed by surgery for thoracic esophageal cancer potentially or actually involving adjacent organs. *Dis Esophagus* 2001;14:197-201.
- 42 Fujiwara H, Terashima M, Irinoda T, Takagane A, Abe K, Kashiwaba M, Oyama K, Takahashi M, Maesawa C, Saito K, Takechi T, Fukushima M. Quantitative measurement of thymidylate synthase and dihydropyrimidine dehydrogenase mRNA level in gastric cancer by real-time RT-PCR. *Jpn J Cancer Res* 2002;93:1342-1350.
- 43 Tarusawa M, Yashima A, Endo M, Maesawa C. Quantitative assessment of minimal residual disease in childhood lymphoid malignancies using an allele-specific oligonucleotide real-time quantitative polymerase chain reaction. *Int J Hematol* 2002;75:166-173.
- 44 Uchiyama M, Maesawa C, Yashima A, Itabashi T, Satoh T, Tarusawa M, Endo M, Takahashi Y, Sasaki S, Tsuchiya S, Ishida Y, Masuda T. Development of immunoglobulin variable heavy chain gene consensus probes with conjugated 3' minor groove binder groups for monitoring minimal residual disease in childhood acute lymphoblastic leukaemia. *J Clin Pathol* 2003;56:952-955.
- 45 ○Akiyama Y, Maesawa C, Ogasawara S, Terashima M, Masuda T. Cell-type-specific repression of the maspin gene is disrupted frequently by demethylation at the promoter

- region in gastric intestinal metaplasia and cancer cells. *Am J Pathol* 2003;163:1911-1919.
- 46 Uchiyama M, Maesawa C, Yashima A, Tarusawa M, Satoh T, Ishida Y, Ito S, Murai K, Enomoto S, Utsugisawa T, Motoyoshi K, Masuda T. Development of consensus fluorogenically labeled probes of the immunoglobulin heavy-chain gene for detecting minimal residual disease in B-cell non-Hodgkin lymphomas. *Cancer Sci* 2003;94:877-885.
- 47 Uchiyama M, Maesawa C, Yashima A, Itabashi T, Ishida Y, Masuda T. Consensus primers for detecting monoclonal immunoglobulin heavy chain rearrangement in B cell lymphomas. *J Clin Pathol* 2003;56:778-779.
- 48 Yashima A, Maesawa C, Uchiyama M, Tarusawa M, Satoh T, Satoh M, Enomoto S, Sugawara K, Numaoka H, Murai K, Utsugisawa T, Ishida Y, Masuda T. Quantitative assessment of contaminating tumor cells in autologous peripheral blood stem cells of B-cell non-Hodgkin lymphomas using immunoglobulin heavy chain gene allele-specific oligonucleotide real-time quantitative-polymerase chain reaction. *Leuk Res* 2003;27:925-934.
- 49 ○Maesawa C, Inaba T, Sato H, Iijima S, Ishida K, Terashima M, Sato R, Suzuki M, Yashima A, Ogasawara S, Oikawa H, Sato N, Saito K, Masuda T. A rapid biosensor chip assay for measuring of telomerase activity using surface plasmon resonance. *Nucleic Acids Res* 2003;31:E4-4.
- 50 Nakatani K, Hagihara S, Sando S, Sakamoto S, Yamaguchi K, Maesawa C, Saito I. Induction of a remarkable conformational change in a human telomeric sequence by the binding of naphthyridine dimer: inhibition of the elongation of a telomeric repeat by telomerase. *J Am Chem Soc* 2003;125:662-666.
- 51 ○Sato R, Maesawa C, Fujisawa K, Wada K, Oikawa K, Takikawa Y, Suzuki K, Oikawa H, Ishikawa K, Masuda T. Prevention of critical telomere shortening by oestradiol in human normal hepatic cultured cells and carbon tetrachloride induced rat liver fibrosis. *Gut* 2004;53:1001-1009.
- 52 Wada K, Maesawa C, Akasaka T, Masuda T. Aberrant expression of the maspin gene associated with epigenetic modification in melanoma cells. *J Invest Dermatol* 2004;122:805-811.
- 53 Oyama K, Terashima M, Takagane A, Maesawa C. Prognostic significance of peritoneal minimal residual disease in gastric cancer detected by reverse transcription-polymerase chain reaction. *Br J Surg* 2004;91:435-443.
- 54 ○Uchiyama M, Maesawa C, Yashima-Abo A, Tarusawa M, Satoh M, Satoh T, Ishida Y, Ito S, Murai K, Enomoto S, Utsugisawa T, Masuda T. Short consensus probes with 3'-minor groove binder of the immunoglobulin heavy-chain gene for real-time quantitative PCR in B-cell non-Hodgkin lymphomas. *Lab Invest* 2004;84:932-936.

- 55 Akiyama Y, Maesawa C, Wada K, Fujisawa K, Itabashi T, Noda Y, Honda T, Sato N, Ishida K, Takagane A, Saito K, Masuda T. Human PinX1, a potent telomerase inhibitor, is not involved in human gastrointestinal tract carcinoma. *Oncol Rep* 2004;11:871-874.
- 56 Itabashi T, Maesawa C, Uchiyama M, Higuchi T, Masuda T. Quantitative detection of mutant alleles of the K-ras gene with minor groove binder-conjugated fluorogenic DNA probes. *Int J Oncol* 2004;24:687-696.
- 57 Ogasawara S, Maesawa C, Yamamoto M, Akiyama Y, Wada K, Fujisawa K, Higuchi T, Tomisawa Y, Sato N, Endo S, Saito K, Masuda T. Disruption of cell-type-specific methylation at the Maspin gene promoter is frequently involved in undifferentiated thyroid cancers. *Oncogene* 2004;23:1117-1124.
- 58 Satoh H, Nakamura M, Satoh M, Nakajima T, Izumoto H, Maesawa C, Kawazoe K, Masuda T, Hiramori K. Expression and localization of tumour necrosis factor-alpha and its converting enzyme in human abdominal aortic aneurysm. *Clin Sci (Lond)* 2004;106:301-306.
- 59 Maesawa C, Oikawa K, Fujiwara-Akita H, Masuda T. Biacore's SPR technology for the development of telomere/telomerase-targeted therapies. *Biacore J* 2004;1:7-9.
- 60 Terashima M, Maesawa C, Oyama K, Ohtani S, Akiyama Y, Ogasawara S, Takagane A, Saito K, Masuda T, Kanzaki N, Matsuyama S, Hoshino Y, Kogure M, Gotoh M, Shirane M, Mori K. Gene expression profiles in human gastric cancer: expression of maspin correlates with lymph node metastasis. *Br J Cancer* 2005;92:1130-1136.
- 61 Iwaya T, Maesawa C, Kimura T, Ogasawara S, Ikeda K, Kimura Y, Noda Y, Ishida K, Sato N, Saito K, Masuda T. Infrequent mutation of the human envoplakin gene is closely linked to the tylosis oesophageal cancer locus in sporadic oesophageal squamous cell carcinomas. *Oncol Rep* 2005;13:703-707.
- 62 Fujiwara-Akita H, Maesawa C, Honda T, Kobayashi S, Masuda T. Expression of human telomerase reverse transcriptase splice variants is well correlated with low telomerase activity in osteosarcoma cell lines. *Int J Oncol* 2005;26:1009-1016.
- 63 Uchiyama M, Maesawa C, Yashima-Abo A, Tarusawa M, Endo M, Sugawara W, Chida S, Onodera S, Tsukushi Y, Ishida Y, Tsuchiya S, Masuda T. Consensus JH gene probes with conjugated 3'-minor groove binder for monitoring minimal residual disease in acute lymphoblastic leukemia. *J Mol Diagn* 2005;7:121-126.
- 64 Fujisawa K, Maesawa C, Sato R, Wada K, Ogasawara S, Akiyama Y, Takeda M, Fujita T, Otsuka K, Higuchi T, Suzuki K, Saito K, Masuda T. Epigenetic status and aberrant expression of the maspin gene in human hepato-biliary tract carcinomas. *Lab Invest* 2005;85:214-224.
- 65 Murai S, Maesawa C, Masuda T, Sugiyama T. Aberrant maspin expression in human

- endometrial cancer. *Cancer Sci* 2006;97:883-888.
- 66 Kimura T, Maesawa C, Ikeda K, Wakabayashi G, Masuda T. Mutations of the epidermal growth factor receptor gene in gastrointestinal tract tumor cell lines. *Oncol Rep* 2006;15:1205-1210.
- 67 Fujita T, Maesawa C, Oikawa K, Nitta H, Wakabayashi G, Masuda T. Interferon-gamma down-regulates expression of tumor necrosis factor-alpha converting enzyme/a disintegrin and metalloproteinase 17 in activated hepatic stellate cells of rats. *Int J Mol Med* 2006;17:605-616.
- 68 Maesawa C, Ogasawara S, Yashima-Abo A, Kimura T, Kotani K, Masuda S, Nagata Y, Iwaya T, Suzuki K, Oyake T, Akiyama Y, Kawamura H, Masuda T. Aberrant maspin expression in gallbladder epithelium is associated with intestinal metaplasia in patients with cholelithiasis. *J Clin Pathol* 2006;59:328-330.
- 69 Satoh M, Shimoda Y, Maesawa C, Akatsu T, Ishikawa Y, Minami Y, Hiramori K, Nakamura M. Activated toll-like receptor 4 in monocytes is associated with heart failure after acute myocardial infarction. *Int J Cardiol* 2006;109:226-234.
- 70 Yamashina M, Kaneko Y, Maesawa C, Kajiwaru T, Ishii M, Fujiwara F, Taneichi H, Takebe N, Ishida W, Takahashi K, Masuda T, Satoh J. Association of TNF-alpha gene promoter C-857T polymorphism with higher serum LDL cholesterol levels and carotid plaque formation in Japanese patients with type 2 diabetes. *Tohoku J Exp Med* 2007;211:251-258.
- 71 Tokita T, Maesawa C, Kimura T, Kotani K, Takahashi K, Akasaka T, Masuda T. Methylation status of the SOCS3 gene in human malignant melanomas. *Int J Oncol* 2007;30:689-694.
- 72 Izutsu N, Maesawa C, Shibazaki M, Oikawa H, Shoji T, Sugiyama T, Masuda T. Epigenetic modification is involved in aberrant expression of class III beta-tubulin, TUBB3, in ovarian cancer cells. *Int J Oncol* 2008;32:1227-1235.
- 73 Itabashi H, Maesawa C, Oikawa H, Kotani K, Sakurai E, Kato K, Komatsu H, Nitta H, Kawamura H, Wakabayashi G, Masuda T. Angiotensin II and epidermal growth factor receptor cross-talk mediated by a disintegrin and metalloprotease accelerates tumor cell proliferation of hepatocellular carcinoma cell lines. *Hepatol Res* 2008;38:601-613.
- 74 Yamauchi K, Piao HM, Nakadate T, Shikanai T, Nakamura Y, Niisato M, Ito H, Mouri T, Kobayashi H, Maesawa C, Sawai T, Ohtsu H, Inoue H. Progress in allergy signal research on mast cells: the role of histamine in goblet cell hyperplasia in allergic airway inflammation - a study using the Hdc knockout mouse. *J Pharmacol Sci* 2008;106:354-360.
- 75 Mitomo S, Maesawa C, Ogasawara S, Iwaya T, Shibazaki M, Yashima-Abo A, Kotani K, Oikawa H, Sakurai E, Izutsu N, Kato K, Komatsu H, Ikeda K, Wakabayashi G, Masuda T. Downregulation of miR-138 is associated with overexpression of human telomerase reverse transcriptase protein in human anaplastic thyroid carcinoma cell lines. *Cancer Sci*

2008;99:280-286.

- 76 Yamauchi K, Piao HM, Nakadate T, Shikanai T, Nakamura Y, Ito H, Mouri T, Kobayashi H, Maesawa C, Sawai T, Ohtsu H, Inoue H. Enhanced Goblet Cell Hyperplasia in HDC Knockout Mice with Allergic Airway Inflammation. *Allergol Int* 2009;58:125-134.
- 77 ○Akasaka K, Maesawa C, Shibazaki M, Maeda F, Takahashi K, Akasaka T, Masuda T. Loss of class III beta-tubulin Induced by histone deacetylation is associated with chemosensitivity to paclitaxel in malignant melanoma cells. *J Invest Dermatol* 2009. in press (Epub ahead of print)
- 78 Nagata Y, Maesawa C, Tada H, Takikawa Y, Yashima-Abo A, Masuda T. Differential microRNA expression between bone marrow side population cells and hepatocytes in adult mice. *Int J Mol Med*, in press.
- 79 Kato K, Maesawa C, Itabashi T, Fujisawa K, Otsuka K, Kanno S, Tada H, Tatemichi Y, Kotani K, Oikawa H, Sugai T, Wakabayashi G, Masuda T. DNA hypomethylation at the CpG island is involved in aberrant expression of the L1 cell adhesion molecule gene in colorectal cancer. *Int J Oncol*, in press.
- 80 Minami Y, Satoh M, Maesawa C, Takahasi Y, Tabuchi T, Itoh T, Nakamura M. Effect of atorvastatin on microRNA221/222 expression from patients with coronary artery disease. *Eur J Clin Invest*, in press.

Case Report

- 1 Maesawa C, Tamura G, Sawada H, Kamioki S, Nakajima Y, Satodate R. Angiomyolipoma arising in the colon. *Am J Gastroenterol* 1996;91:1852-1854.
- 2 Iwaya T, Maesawa C, Tamura G, Sato N, Ikeda K, Sasaki A, Othuka K, Ishida K, Saito K, Satodate R. Esophageal carcinosarcoma: a genetic analysis. *Gastroenterology* 1997;113:973-977.
- 3 Kimura Y, Kawamura M, Owada M, Fujiwara T, Maesawa C, Hiramori K. Successful steroid therapy for cefdinir-induced acute tubulointerstitial nephritis with progressive renal failure. *Intern Med* 2001;40:114-117.
- 4 Wada K, Maesawa C, Satoh T, Akasaka T, Masuda T. A case of primary cutaneous CD30+ T-cell lymphoproliferative disorder with features of granulomatous slack skin disease. *Br J Dermatol* 2002;147:998-1002.
- 5 Iijima S, Maesawa C, Sato N, Ikeda K, Inaba T, Akiyama Y, Ishida K, Saito K, Masuda T. Gastrointestinal stromal tumour of the oesophagus: significance of immunohistochemical and genetic analyses of the c-kit gene. *Eur J Gastroenterol Hepatol* 2002;14:445-448.

- 6 Maesawa C, Iijima S, Sato N, Yoshinori N, Suzuki M, Tarusawa M, Ishida K, Tamura G, Saito K, Masuda T. Esophageal extraskeletal Ewing's sarcoma. *Hum Pathol* 2002;33:130-132.
- 7 Sato A, Imaizumi M, Chikaoka S, Niizuma H, Hoshi Y, Takeyama J, Fujii K, Nishio T, Watanabe M, Maesawa C, Hayashi Y, Inuma K. Acute renal failure due to leukemic cell infiltration followed by relapse at multiple extramedullary sites in a child with acute lymphoblastic leukemia. *Leuk Lymphoma* 2004;45:825-828.
- 8 Oikawa H, Maesawa C, Sato R, Oikawa K, Yamada H, Oriso S, Ono S, Yashima-Abo A, Kotani K, Suzuki K, Masuda T. Liver cirrhosis induced by long-term administration of a daily low dose of amiodarone: a case report. *World J Gastroenterol* 2005;11:5394-5397.
- 9 Yashima-Abo A, Satoh T, Abo T, Aoki Y, Kowata S, Ito S, Ishida Y, Fujiwara H, Maesawa C, Masuda T. Distinguishing between proliferating nodal lymphoid blasts in chronic myelogenous leukemia and non-Hodgkin lymphoma: report of three cases and detection of a bcr/abl fusion signal by single-cell analysis. *Pathol Int* 2005;55:273-279.
- 10 Nakamura Y, Sato A, Takahashi S, Kanazawa T, Ito S, Maesawa C, Masuda T, Inoue Y. Primary pericardial mesothelioma presenting as constrictive pericarditis. *Respiratory Med. Extra* 2005;1:107-109.
- 11 Iwaya T, Maesawa C, Uesugi N, Kimura T, Ikeda K, Kimura Y, Mitomo S, Ishida K, Sato N, Wakabayashi G. Coexistence of esophageal superficial carcinoma and multiple leiomyomas: A case report. *World J Gastroenterol* 2006;12:4588-4592.
- 12 Iwaya T, Maesawa C, Uesugi N, Kimura T, Ogasawara S, Ikeda K, Kimura Y, Mitomo S, Ishida K, Sato N, Saito K, Masuda T. True carcinosarcoma of the esophagus. *Dis Esophagus* 2006;19:48-52.
- 13 Nishimura K, Satoh T, Maesawa C, Ishijima K, Sato H. Giant cell tumor of the larynx: a case report and review of the literature. *Am J Otolaryngol* 2007;28:436-440.